

Childhood Immunization Schedule

Growing Your Family?

To protect your baby from whooping cough, RSV, and flu, pregnant people need a Tdap vaccine between 27-36 weeks, a RSV vaccine between 32-36 weeks, and a flu vaccine at any time during pregnancy.

BIRTH

Birth

- Hepatitis B #1

1-2 MONTHS

1-2 months

- Hepatitis B #2
- Diphtheria, tetanus, and pertussis (whooping cough) (DTaP) #1
- Haemophilus influenzae type b (Hib) #1
- Polio #1
- Pneumonia #1
- Rotavirus #1

4 MONTHS

4 months

- Diphtheria, tetanus, and pertussis (whooping cough) (DTaP) #2
- Haemophilus influenzae type b (Hib) #2
- Polio #2
- Pneumonia #2
- Rotavirus #2

Between birth-8 months

- RSV (depending on maternal RSV vaccination status)

6 MONTHS

6 months

- Flu (depending on time of year) #1
- Pneumonia #3
- Diphtheria, tetanus, and pertussis (whooping cough) (DTaP) #3
- Rotavirus (depending on manufacturer) #3
- Haemophilus influenzae type b (Hib) #3 (depending on manufacturer)

12 MONTHS

12-15 months

- Haemophilus influenzae type b (Hib) #4 (depending on manufacturer)
- Measles, mumps, and rubella (German measles) (MMR) #1
- Pneumonia #4
- Chickenpox #1

15 MONTHS

15 months-18 months

- Diphtheria, tetanus, and pertussis (whooping cough) (DTaP) #4

18 MONTHS

Between 6-18 months

- Hepatitis B #3
- Polio #3
- Second dose of flu

Between 12-23 months

- Hepatitis A (2 doses separated by 6 months)

FLU VACCINE

The CDC recommends children 6 months and older get a flu vaccine every year.

Kids 2 and older may be eligible to get a mist vaccine rather than injection.

Children under 9 who have not gotten the vaccine before should plan on two doses, four weeks apart.

COVID-19 VACCINE

All eligible individuals, including children, should receive the recommended doses of COVID-19 vaccine.

4-6 YEARS

4-6 years

- Diphtheria, tetanus, and pertussis (whooping cough) (DTaP) #5
- Measles, mumps, and rubella (German measles) (MMR) #2
- Polio #4
- Chickenpox #2

11-12 YEARS

11-12 years

- Tetanus, diphtheria, and pertussis (whooping cough) (Tdap) #1
- HPV #1 and #2
- Meningitis ACWY #1

16-18 YEARS

16-18 years

- Meningitis ACWY #2
- Meningitis B (2 or 3 doses depending on manufacturer)